

# CUT&RUN library preparation

 Paolo Emanuele Forni

Updated date: Jun 20, 2023

 An abbreviated version of this protocol was published in eLIFE

Sociosexual behavior requires both activating and repressive roles of *Tfap2e*/AP-2 $\epsilon$  in vomeronasal sensory neurons

DOI: 10.7554/eLife.77259

## Detailed protocol

Fragment analysis of prepared CUT&RUN libraries was performed as a fee-for-service via Cornell University's Genomics Core facility. DNA library quality was assessed by qualitatively visualizing fragment traces for fragments of the expected size (200-1000bp). Traces displaying empty adaptors (130bp) or free floating primers (65bp) were subjected to an additional SPRI bead cleanup to exclude unwanted fragments.

**How to cite:** (Readers should cite both the Bio-protocol preprint and the original research article where this protocol was used)

1. Forni, P. E.(2023). CUT&RUN library preparation. Bio-protocol Preprint. [bio-protocol.org/prep2341](https://bio-protocol.org/prep2341).
2. Sociosexual behavior requires both activating and repressive roles of *Tfap2e*/AP-2 $\epsilon$  in vomeronasal sensory neurons. eLIFE. DOI: [10.7554/eLife.77259](https://doi.org/10.7554/eLife.77259)

**Copyright:** © 2023 The Author(s); This is an open-access article under the CC BY-NC license (<https://creativecommons.org/licenses/by-nc/4.0/>).